



BROMOTRIL T

Reg.No. L6653 Act / Wet 36/1947
N-AR 0870

A selective suspension concentrate herbicide for the post-emergence control of certain broadleaf weeds in maize, grain sorghum, sugarcane, sweetcorn and fodder sorghum.

'n Selektiewe na-opkom suspensie konsentraat onkruidodder vir die beheer van sekere breëblaaronkruid in mielies, graansorghum, suikerriet, suikermielies en voersorghum.

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE

Bromoxynil (Nitrile).....150 g / l.....Bromoksiniel (Nitriël)
(Heptanoate and octanoate esters) (Heptanoaat en oktanoaat esters)

Terbutylazine (Triazine).....333 g / l.....Terbutilasien (Triasion)

NETT CONTENTS.....NETTO INHOUD

REGISTRATION HOLDER / REGISTRASIEHOUER:

MAKHTESHIM – AGAN SA (PTY) LTD

Reg No. 1992 / 001741 / 07

21 Viben Street

BRACKENFELL

7560

Tel No (021) 982 1460

Batch number.....Lotnommer

Date of Manufacture.....Datum van Vervaardiging



**HARMFULL
SKADELIK**



WARNINGS

A minimum of 14 days should be allowed between application and grazing or feeding:

- Handle with care
- Poisonous if swallowed, inhaled or absorbed through the skin.
- Irritating to eyes and may cause sensitisation by skin contact.
- Toxic to fish, bees and wildlife.
- Store in a cool, dry place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.
- May cause harm to the unborn child.

Aerial application:

Aerial application of **BROMOTRIL T** may only be done by a registered Aerial Application Operator using a correctly calibrated, registered aircraft according to the instructions of SABS Code 0118 (Aerial Application of Agricultural Pesticides). Ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria be met:

- **Volume:** A spray mixture volume of 30 (at least) to 35 (recommended) litres per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.
- **Droplet coverage:** 35 to 45 droplets per cm² must be recovered at the target area.
- **Droplet size:** A droplet spectrum with a VMD of 350 microns is recommended. Limit the production of fine droplets less than 150 microns (high drift and evaporation potential) to a minimum.
- **Flying height:** Maintain the height of the spray boom at 3 to 4 metres above the target. Do not spray when aircraft dives, is in a climb or when banking
- Use suitable atomising equipment that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product. The spraying system must produce a droplet spectrum with the lowest possible Relative Span.
- Position all the atomisers within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- The difference in temperature between the wet and dry bulb thermometers, of a whirling hygrometer, should not exceed 8°C.
- Stop spraying if the wind speed exceeds 15 km/h.
- Stop spraying under turbulent, unstable and dry conditions during the heat of the day.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - a) reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage).
 - b) damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that the Aerial Spray Operator knows exactly which fields to spray.
- Obtain an assurance from the Aerial Spray Operator that the above requirements will be met and that relevant data will be compiled in a logbook and kept for future reference.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by the factors such as abnormal climatic and storage conditions, the quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of weeds against the remedy concerned as well as by the method time and accuracy of application. The registration holder further more does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Avoid inhalation of fumes and spray mist.
- Wear protective clothing, overalls, boots, and rubber gloves while mixing or applying the product.
- Wash with soap and water after application.
- Remove and wash contaminated clothing after a days work.
- In case of eye contact wash out eyes with clean running water for at least 15 minutes.
- Do not eat, drink or smoke whilst mixing or applying of before hands and face have been washed and clothes have been changed.
- Avoid contamination of food, eating utensils, feed and drinking water.
- Avoid drift of spray mist to other crops, grazing, rivers, dams or any other area that is not under treatment.
- Do not mix and load within 15 m from boreholes, rivers, streams or dams.
- Do not apply within 15 m from boreholes, rivers and streams.
- Do not apply within 60 m from dams.

- Clean applicator thoroughly after use. Dispose of wash water where it will not contaminate the environment.
- Invert the empty container over the spray tank and allow to drain for a minimum of 30 seconds after the flow has slowed down to a drip, thereafter rinse the container three times with a volume of water equal to at least 10% of that of the container. Add the rinsings to the contents of the spray tank. Dispose of the container by making holes in it, flattening it and burying it. Do not use the container for any other purpose.

USE RESTRICTIONS

To prevent damage to follow on crops, the following waiting periods after the application of BROMOTRIL T should be adhered to:

Maize and Sugarcane	None
All other crops	4 Months

EXCEPT

Where the clay content of the soil is less than 20% the organic material content is less than 1,0% and the soil pH is more than 7 the waiting period for all other crops changes to 12 months.

DIRECTIONS FOR USE USE ONLY AS DIRECTED

BROMOTRIL-T is compatible with Atrazine SC formulations and the mixture can be used if a longer residual action is needed on susceptible broadleaf weeds.

MIXING INSTRUCTIONS

SHAKE THE BROMOTRIL-T CONTAINER THOROUGHLY BEFORE USE.
REPLACE AND TIGHTEN THE CAP AFTER USE.

Half fill the spray tank with clean water. Mix the required volume of BROMOTRIL-T separately in water and add to the spray tank while agitating the spray mixture. Fill the spray tank with water while agitation is maintained. If BROMOTRIL-T is used in a tank mixture with Atrazine SC formulations the BROMOTRIL-T should be added first into the half filled spray tank and the Atrazine SC thereafter. Fill the tank with water while agitation is maintained. The spray mixture must be agitated continuously before and during application. The spray mixture should never be allowed to stand overnight.

APPLICATION

To achieve effective weed control, thorough wetting of the target weeds is essential.

GROUND APPLICATION

A tractor mounted sprayer filled with a conventional boom and flat nozzles can be used for the application of BROMOTRIL-T. The pressure during spraying should not exceed 300 kPa. To ensure thorough wetting of the target weeds at least 250-300 l water per ha should be used.

If the crop is higher than 400 mm and the foliage is very dense, drop arms can be used to ensure thorough wetting of the target weeds.

AERIAL APPLICATION

BROMOTRIL-T can be applied aerially provided that the spray mixture is applied evenly over the target area and that the loss of chemical is limited to the minimum during application. To achieve this, the following conditions must be met:

- A conventional spray boom fitted with flat fan nozzles eg. SS 6515 or similar should be used.
- A minimum spray volume of 30 l but preferably 35 l per ha must be applied.
- A minimum of 30-50 droplets per square centimetre should be recovered on the target area.
- The droplet spectrum must have a VMD of 350 micron.
- Maintain an altitude of 3 m above the target area at wind speeds of 0-8 km/h and 2 m above the target area at wind speeds of 8-15 km/h. Stop spraying when wind speed exceeds 15 km/h.
- The difference between the readings of the wet and dry bulbs as determined by a whirling hygrometer should not exceed 8°C
- Do not spray during the heat of the day.

It is essential to get assurance of the spray operator that the above mentioned conditions will be met.

APPLICATION RATES

CROP	DOSAGE RATE per ha	REMARKS
MAIZE AND GRAIN SORGHUM	Ground application: 1,25 l Aerial application 1,5 l	Apply post-emergence when most of the weeds have germinated but before they have developed beyond the 4-leaf stage except if stated otherwise on the lists below.
	Ground application: 1,5 l Aerial application: 1,8 l	Apply post-emergence when most of the weeds have germinated but before they have developed beyond the 6-leaf stage except if stated otherwise on the lists below.
	Ground application: 1,25 l BROMOTRIL-T + 1,0-2,0 l Atrazine SC Aerial application: 1,5 l BROMOTRIL-T + 1,0-2,0 l Atrazine SC	If longer residual action is needed, tank mixes with Atrazine SC can be made as follows: Apply post-emergence when most of the weeds have germinated but before they have developed beyond the 4-leaf stage , except if stated otherwise on the lists below.
	Ground application: 1,5 l BROMOTRIL-T + 1,0-2,0 l Atrazine SC	Apply post-emergence when most of the weeds have germinated but before they have developed beyond the 6-leaf stage , except if stated otherwise on the lists below.
SWEETCORN AND FODDER SORGHUM	Ground application: 1,25 l	Apply post-emergence when most of the weeds have germinated, but before they have developed beyond the 4-leaf stage , except if stated otherwise on the lists below.
	Ground application: 1,5 l	Apply post-emergence when most of the weeds have germinated, but before they have developed beyond the 6-leaf stage , except if stated otherwise on the lists below.
SUGARCANE (Plant and ratoon cane)	Ground application: 2,0 l	Apply post-emergence when most of the weeds have germinated but before they have developed beyond the 6-leaf stage , except if stated otherwise on the lists below.

IMPORTANT INFORMATION

1. If the crop is subjected to stress conditions eg. High temperatures, drought, low temperatures, nutrient deficiencies, insect or nematode damage, etc. during or shortly after the application of BROMOTRIL-T, chlorosis or leaf scorch might appear on the crop. Crop plants usually recover completely with no detrimental effect on yield.

If maize plants are subjected to streak disease serious damage might occur after a BROMOTRIL-T application.

2. If weeds have developed beyond the growth stage as indicated on the label at application of BROMOTRIL-T, poor weed control may result.
3. To obtain optimum weed control, weeds should be growing actively at the time of application. Any stress condition will be detrimental to weed control.
4. Thorough wetting of weeds should be ensured to obtain good weed control. Drop arms can be used if the crop is too high or foliage growth is too dense to ensure thorough wetting.
5. BROMOTRIL-T controls susceptible annual broad leaf weeds only. To control annual grasses a suitable pre-emergence grass herbicide should be used.
6. When Atrazine SC is used in a tank mix with BROMOTRIL-T, the Atrazine SC label should be studied carefully and not should also be taken regarding the waiting periods.
7. No adjuvants must be used with BROMOTRIL-T as they can cause damage to the crop or even be detrimental on the effectivity.
8. Do not mix BROMOTRIL-T with any other compound, except Atrazine SC, before the supplier has been consulted.

WAARSKUWINGS

'n Minimum periode van 14 dae moet toegelaat word tussen toediening en beweiding of voer.

- Hanteer versigtig.
- Giftig indien ingesluk, ingeasem of deur die vel geabsorbeer word.
- Irreterend vir oë en mag sensitisering deur velkontak veroorsaak.
- Giftig vir visse, bye en die natuurlewe.
- Berg in 'n koel plek, weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Moontlike risiko dat dit skadelik kan wees teenoor 'n ongebore kind.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registratiewaerder nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimats- en bergingstoestande, kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die onkruid teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiewaerder nie verantwoordelikeheid vir skade aan gewasse, plantegroei, die omgewing of vir die nadelige effek op mens en dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie krans die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS

- Moet nie dampe of sproeiwater inasem nie.
- Dra beskermende oorklere (oorpak, stewels en rubberhandskoene) wanneer die produk vermeng of toegedien word. Was met water en seep na toediening.
- Trek besoedelde klere uit en was daaglik.
- In die geval van oogkontak spoel met lopende water uit vir ten minste 15 minute.
- Moet nie eet, drink of rook tydens vermenging of toediening of voordat hande en gesig gewas is en skoon klere aangetrek is nie.
- Verhoed die besoedeling van voedsel, eetgerei, voer en drinkwater.
- Moet nie meng en laai nader as 15 m vanaf boorgate, riviere, strome en damme nie.
- Moet nie nader as 15 m vanaf boorgate, riviere, en strome toedien nie.
- Moet nie nader as 60 m vanaf damme toedien nie.
- Dra sorg dat geen terugsyfering deur besproeiingstelsels na boorgate en damme plaasvind nie.
- Maak die spuittoerusting deeglik skoon na toediening en gooi die waswater weg waar dit nie die omgewing kan besoedel nie.
- Keer leë houer om oor die spuittenk en dreineer vir tenminste 30 sekondes nadat die vloeitot 'n gedrup verminder het. Spoel daarna die houer drie keer uit met 'n volume water wat gelykstaande aan minstens 10% van die volume van die leë houer is. Gooi die spoelwater by die inhoud van die spuittenk. Vernietig die leë houer deur gate daarin te maak, plat te druk en dit te begrawe.
- Moet leë houer vir geen ander doel gebruik nie.

GEBRUIKSBEPERKINGS

Om skade aan gevoelige opvolggewasse te beperk, moet die volgende wagperiodes na toediening van BROMOTRIL-T gehandhaaf word:

Mielies en Suikerriet
Alle ander gewasse

Geen
4 Maande

BEHALWE

Waar die klei-inhoud van die grond minder as 20% is, organiese materiaalinhoud minder as 1% is en die pH van die grond meer as 7, is verander die wagperiode vir alle gevoelige gewasse na 12 maande.

GEBRUIKSAANWYSINGS GEBRUK SLEGS SOOS AANGEDUI

VERENIGBAARHEID

BROMOTRIL-T is verenigbaar met Atrasien SC formulasies en die mengsel kan gebruik word om langer nawerking op gevoelige breëblaaronkruid te bewerkstellig.

VERMENING

SKUD DIE KAN DEEGLIK VOOR GEBRUIK. DRAAI DIE SKROEFPROP VAS NA GEBRUIK.

Vul die spuitnek half vol met skoon water. Vermeng die benodigde hoeveelheid BROMOTRIL-T apart in water en voeg in spuitnek terwyl gedurigdeur geroer word. Vul die spuitnek tot op die verlangde volume met water terwyl geroer word.

Indien BROMOTRIL-T met 'n Atrasien SC formulasie in 'n tenkmengsel gebruik word moet die BROMOTRIL-T eerste in die halfvol spuitnek gevoeg word terwyl geroer word. Die Atrasien SC formulasie word daarna in die spuitnek gevoeg terwyl geroer word. Die oorlywende volume water word dan in die spuitenknog gevoeg.

Die spuitmengsel moet voor en tydens toediening gedurigdeur geroer word. Die spuitmengsel moet nie oornag laat staan word nie.

TOEDIENING

'n Deeglike benatting van onkruid is noodsaaklik vir doeltreffende beheer.

GRONDTOEDIENING

'n Trekkerspuit toegerus met 'n konvensionele spuitbalk en plat waaierspuitpunte, kan vir die toediening van BROMOTRIL-T gebruik word. Die spuitdruk moet nie 300 kPa oorskry nie. Om deeglike benatting van die onkruid te verkry moet minstens 250-300 l water per ha gebruik word.

Indien die gewas hoër as 400 mm is en ruig groei, kan van valarms gebruik gemaak word om 'n deeglike bedekking van onkruid te verkry.

Lugtoediening:

BROMOTRIL T kan slegs deur 'n geregistreerde Lugbespuitingsperateur met 'n korrek gekalibreerde, geregistreerde vliegtuig volgens die instruksies van SABS Kode 0118 (Aerial Application of Agricultural Pesticides) uit die lug bespuit word. Verseker dat die spuitmengsel eweredig oor die teikenarea versprei word, en die verlies aan spuitmengsel tydens toediening tot 'n minimum beperk word. Dit is daarom belangrik om aan die volgende vereistes te voldoen:

- Volume: 'n Spuitmengsel volume van 30 (minimum) tot 35 (aanbevole volume) liter per ha word aanbeveel. Hierdie produk is nie teen 'n verlaagde volume getoets nie. Die registrasiehouer kan nie effektiwiteit waarborg, of verantwoordelik gehou word vir enige nadelige effekte indien hierdie produk teen 'n laer volume, as hierbo aanbeveel, toegedien word nie.
- Druppel bedekking: 35 tot 45 druppels per cm² moet op die teikenarea herwin word.
- Druppelgrootte: 'n Druppelspektrum met 'n VMD van 350 mikrons word aanbeveel. Beperk die produksie van druppels kleiner as 150 mikrons (hoë drywing en verdampingspotensiaal) tot 'n minimum.
- Vlieghoogte: Handhaaf die hoogte van die spuitbalk bo die teiken op 3 tot 4 meter. Moet nie spuit wanneer die vliegtuig duik nie, uitklim of draai nie.
- Gebruik geskikte atomiseringsapparaat wat die vereiste druppelgrootte en bedekking sal produseer, maar die minste verlies van produk verseker. Die spuitstelsel moet 'n druppelspektrum met die kleinste moontlike Relatiewe Span produseer.
- Plaas al die atomiseerders in die binnste 60 tot 75 % van die vlerkspan om te verhoed dat druppels binne-in die vlerkpuntvorteks beweeg.
- Die verskil in temperatuur tussen die nat- en droëboltermometer van 'n swaaihigrometer, moet nie 8°C oorskry nie.
- Stop bespuiting indien die windspoed 15 km/h oorskry.
- Stop bespuiting tydens turbulente, onstabiele en droë toestande gedurende die hitte van die dag.
- Bespuiting onder temperatuur inversie toestande (deur bo of binne die inversie laag te spuit) en/of hoë lugvog toestande (relatiewe humiditeit 80% en meer) mag tot volgende probleme aanleiding gee:
 - a) verlaagde effektiwiteit aangesien die druppels as 'n wolk in die lug bly hang en moontlik verdamp (onvoldoende bedekking op teiken).
 - b) skade aan nie-teiken gewasse of sensitiewe areas as gevolg van wegdrywing van die spuitwolk na nie-teiken area.
- Verseker dat die Lugbespuitingsoperateur presies weet watter lande bespuit moet word.
- Dit is noodsaaklik om 'n versekering van die Lugbespuitingsoperateur te verkry dat aan al die bogenoemde vereistes voldoen sal word en dat data van belang in 'n logboek saamgevat is vir toekomstige verwysing.

TOEDIENINGSHOEVEELHEDE

GEWAS	TOEDIENINGS- HOEVEELHEID per ha	OPMERKINGS
MIELIES EN GRAANSORGHUM	Grondtoediening: 1,25 l Lugtoediening: 1,5 l	Dien na-opkom toe wanneer die meeste onkruid alreeds opgekom het, maar nog nie verder gevorderd is as die 4-blaarstadium nie, behalwe as anders vermeld in die onderstaande tabelle.
	Grondtoediening: 1,5 l Lugtoediening: 1,8 l	Dien na-opkom toe wanneer die meeste onkruid alreeds opgekom het, maar nog nie verder gevorderd is as die 6-blaarstadium nie, behalwe as anders vermeld word in onderstaande tabelle.
	Grondtoediening: 1,25 l BROMOTRIL-T + 1,0-2,0 l Atrasien SC Lugtoediening: 1,5 l BROMOTRIL-T + 1,0-2,0 l Atrasien SC	Indien langer nawerking verlang word, kan die volgende tenkmengsels met Atrasien SC toegedien word:
	Grondtoediening: 1,25 l BROMOTRIL-T + 1,0-2,0 l Atrasien SC Lugtoediening: 1,5 l BROMOTRIL-T + 1,0-2,0 l Atrasien SC	Dien na-opkom toe wanneer die meeste onkruid alreeds opgekom het maar nog nie verder gevorderd is as die 4-blaarstadium nie, behalwe as anders vermeld word in die onderstaande tabelle.
SUIKERMIELIES EN VOERSORGHUM	Grondtoediening: 1,25 l	Dien na-opkom toe wanneer die meeste onkruid alreeds opgekom het maar nog nie verder gevorderd het as die 4-blaarstadium nie, behalwe as anders vermeld in die onderstaande tabelle.
	Grondtoediening: 1,5 l	Dien na-opkom toe wanneer die meeste onkruid alreeds opgekom het maar nog nie verder gevorderd het as die 6-blaarstadium nie, behalwe as anders vermeld in die onderstaande tabelle.
SUIKERRIET (Plant en oorsaak riet)	Grondtoediening: 2,0 l	Dien na-opkom toe wanneer die meeste onkruid alreeds opgekom het maar nog nie verder gevorderd het as die 6-blaarstadium nie, behalwe as anders vermeld in die onderstaande tabelle.

BELANGRIKE INLIGTING

- Indien die gewas aan stremming onderworpe is as gevolg van hoë temperatuur, droogte, koue, voedingstekorte, insek of aalwurmbesmetting ens. Mag vergelyking

voorkom na toediening van BROMOTRIL-T. Normaalweg herstel die plante volledig en is daar geen invloed op die oesopbrengs nie.

As mielies onderhewig is aan streepsiekte, mag ernstige skade voorkom.

2. Indien onkruid verder ontwikkel het as wat gespesifiseer is, mag dit aanleiding gee tot swak onkruidbeheer.
3. Onkruid moet aktief groei tydens die toediening van BROMOTRIL-T. Indien onkruid tydens toediening onder enige tipe stremming onderworpe is, mag dit tot swak beher van onkruid aanleiding gee.
4. Deeglike benatting van die onkruid met BROMOTRIL-T moet verseker word. Indien die gewas te hoog en te ruig groei om deeglike benatting te verkry, kan van valarms gebruik gemaak word.
5. BROMOTRIL-T beheer net gevoelige eenjarige breëblaaronkruid. Vir die beheer van eenjarige grasse moet van 'n geskikte vooropkom grasdoder gebruik gemaak word.
6. As BROMOTRIL-T saam met Atrasien SC in 'n tenkmengsel toegedien word moet die instruksies op die Atrasien SC etiket noukeurig gelees word en moet daar ook aan die wag periodes van Atrasien SC voldoen word.
7. Geen bevorderingsmiddel moet saam met BROMOTRIL-T gebruik word nie aangesien dit skade aan die gewas kan veroorsaak of selfs die effektiwiteit daarvan kan benadeel.
8. Moet nie BROMOTRIL-T met ander chemikalieë as Atrasien SC formulasies meng voordat die verskaffer nie eers geraadpleeg is nie.

ONKRUIDE WAT DEUR BROMOTRIL-T BEHEER WORD INDIEN TOEDIENING TUSSEN DIE 4-6 BLAARSTADIUM VAN DIE ONKRUIDE TEEN DIE AANBEVOLE TOEDIENINGSHOEVEELHEDE PLAASVIND.

WEEDS CONTROLLED BY BROMOTRIL-T WHEN APPLIED AT THE 4-6 LEAF STAGE OF THE WEEDS AT THE DOSAGE RATES AS RECOMMENDED.

ONKRUIDE / WEEDS	ALGEMENE NAAM	COMMON NAME
<i>Acanthospermum hispidum</i>	Regopsterklits	Upright starbur
<i>Alternanthera pungens</i>	Kakiedubbeltjie	Paperthorn
<i>Amaranthus hybridus</i>	Gewone misbredie	Common pigweed
<i>Amaranthus spinosus</i>	Doringmisbredie	Thorney pigweed
<i>Argemone mexicana</i>	Geelblombloudissel	Yellow flowered Mexican poppy
<i>Bidens pilosa</i>	Gewone knapsekêrel	Common blackjack
<i>Capsella bursa-pastoris</i>	Herderstassie	Shepards purse
<i>Chenopodium album</i>	Withondebossie	White goosefoot
<i>Chenopodium carrinatum</i>	Groenhondebossie	Green goosefoot
<i>Citrullus lanatus</i>	Karkoer	Wild watermelon
<i>Cleome gynandra</i>	Snotterbelletjie	Spider-wisp
<i>Commelina benghalensis</i>	Wandelende jood	Wandering jew
<i>Cosmos bipinnatus</i>	Kosmos	Cosmos
<i>Cucumis myriocarpus</i>	Streepwildekomkommer	Striped wild cucumber
<i>Fallopia convolvulus</i>	Slingerduisendknoop	Climbing knotweed
<i>Flaveria bidentis</i>	Smeltersbossie	Smelter's bush
<i>Galingsoga parviflora</i>	Knopkruid	Gallant soldier
<i>Hibiscus trionum</i>	Terblansbossie	Bladderweed
<i>Ipomoea coscinosperma</i>	-	-
<i>Ipomoea purpurea</i>	Purperwinde	Morning glory
<i>Phytolacca angulata</i>	Wilde-appelliefie	Wild gooseberry
<i>Polygonum aviculare</i>	Voëduisendknoop	Prostrate knotweed
<i>Raphanus repnanistrum</i>	Ramenas	Wild radish
<i>Schkuhria pinnata</i>	Kleinkakaiebos	Dwarf marigold
<i>Sesamum triphyllum</i>	Wildesesam	Wild sesame
<i>Sida cordifolia</i>	Hartblaartaaiman	Flannel weed
<i>Solanum nigrum</i>	Nastergal	Nightshade
<i>Sonchus oleraceus</i>	Sydissel	Sowthistle
<i>Tagetes minuta</i>	Langkakaiebos	Tall Khaki Weed
<i>Xanthium spinosum</i>	Boetebossie	Spiny cocklebur
<i>Xanthium strumarium</i>	Kankerroos	Cocklebur

ONKRUIDE WAT BEHEER WORD DEUR BROMOTRIL-T IN DIE GROEISTADIUMS SOOS HIERONDER AANGEDUI.

WEEDS CONTROLLED BY BROMOTRIL-T AS INDICATED IN THE GROWTH STAGES UNDERNEATH AS INDICATED.

Onkruid / Weed	Algemene Naam	Common name	Dosis / Dosage	
			1,25 l / ha (grond)	1,5 l / ha (Grond)
			1,5 l / ha (lug)	1,8 l / ha (lug)
<i>Amaranthus thunbergii</i>	Rooimisbredie	Red Pigweed	4-blaar / leaf	6-blaar / leaf
<i>Cleome monophylla</i>	Rusperbossie	Spindlepod	-	3-blaar / leaf
<i>Crotalaria sphaerocarpa</i>	Mielie Crotalaria	Wild lucerne	-	50 mm hoog/high
<i>Datura ferox</i>	Grootstinkblaar	Large thorn-apple	6-blaar / leaf	8-blaar / leaf
<i>Datura Stramonium</i>	Olieboom	Common thorn-apple	6-blaar / leaf	8-blaar / leaf
<i>Portulaca oleracea</i>	Porselein	Purslane	-	3-blaar / leaf
<i>Tribulis terrestris</i>	Dubbeltjie	Common dubbetjie	50 mm deursnee / diameter	100 mm deursnee/ diameter