



COTTONEX P

Reg no. L 5868 Act / Wet 36 / 1947

N-AR 0488

<p>A suspension concentrate herbicide for selective pre-emergence control of broadleaf weeds in irrigated cotton when used in combination with a registered grass-killer. It may also be used as a spot emergence directed application.</p>	<p>'n Suspensiekonsentraat onkruidodder vir selektiewe vooropkom beheer van breëblaaronkruid in katoen onder besproeiing wanneer dit in kombinasie met 'n geregistreerde grasodder gebruik word. Dit kan ook as 'n gerigte na-opkombesproeiing toegedien word.</p>
---	--

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE

Prometryn (triazine).....250 g / l.....Prometrien (triasien)

Fluometuron (urea-derivative).....250 g / l.....Fluometuron (urea-derivaat)

NET VOLUME / NETTO VOLUME

.....

REGISTRATION HOLDER / REGISTRASIEHOUER:

MAKHTESHIM-AGAN SA (PTY) LTD

Reg no 1992/001741/07

21 Viben Street

PO Box 498, Brackenfell 7561

Tel no: (021) 982 1460

Batch number

Lotnommer

Date of Manufacture

Datum van vervaardiging



WARNINGS

- Poisonous if swallowed.
- Toxic to fish.
- Store in a cool place.
- Store away from food and feeds.
- Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not guarantee that it will be effective under all conditions. The activity and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of weeds against the remedy, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal, or for lack of performance of the remedy concerned due to failure by the user to follow the label instructions, or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Do not inhale the spray mist.
- Avoid skin contact.
- Wash with soap and water after use.
- Wash contaminated clothing after use.
- Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face.
- Do not mix and load within at least 15 m from boreholes, streams, rivers and dams.
- Do not apply within at least 15 m from boreholes, streams and rivers.
- Do not apply within 60 m from dams.
- Ensure that no back-siphoning to boreholes or dams takes place when product is applied through the irrigation system.
- Avoid drift of spray onto other crops, grazing, rivers, boreholes, dams and areas not under treatment.
- Clean applicator after use. Dispose of rinsate where it will not contaminate crops, grazing, rivers, dams and boreholes.
- Prevent contamination of food, feeds, drinking water, boreholes and eating utensils.
- Rinse the empty container three times with a volume of clean water equal to a minimum of 10% of the container. Add the rinsate to the content of the spray tank before destroying the container.
- Do not use the empty container for any other purpose.

USE RESTRICTIONS

- Do not apply **COTTONEX P** pre-emergence to soils where an attack of damping-off disease is expected, unless effective measures have been taken to combat the disease.
- Do not apply **COTTONEX P** pre-emergence to soils with less than 20% clay content.
- Do not apply **COTTONEX P** pre-emergence to gravelly soils.
- Do not plant any crop other than cotton on lands treated with **COTTONEX P** in the same season.

- Do not use **COTTONEX P** pre-emergence on newly deposited alluvial soils as found in the Uppington and Burgersfort areas and on the banks of the Limpopo river.
- Do not use **COTTONEX P** as a post-emergence directed treatment after it has been applied pre-emergence.
- Do not apply **COTTONEX P** to brackish or alkaline soils.

WEEDS CONTROLLED

The following broadleaf weeds are normally controlled by **COTTONEX P** when used in combination with a registered grass-killer:

<i>Acanthospermum hispidum</i>	Upright starbur
<i>Amaranthus hybridus</i>	Common pigweed
<i>Amaranthus spinosus</i>	Thorney pigweed
<i>Amaranthus thunbergii</i>	Red pigweed
<i>Bidens bipinnata</i>	Spanish blackjack
<i>Bidens pilosa</i>	Blackjack
<i>Flaveria bidentis</i>	Smelter's bush
<i>Galinsoga parviflora</i>	Gallant soldier
<i>Nicandra physaloides</i>	Apple of Peru
<i>Physalis angulata</i>	Wild gooseberry
<i>Portulaca oleracea</i>	Purslane
<i>Richardia brasillensis</i>	Tropical Richardia
<i>Schkuhria pinnata</i>	Dwarf marigold
<i>Sesbania punicea</i>	Red sesbania
<i>Solanum nigrum (complex)</i>	Nightshade
<i>Tagetes minuta</i>	Khaki weed
<i>Zeleya pentandra</i>	Zeleya
Control of the following weeds is variable:	
<i>Cassia italica</i>	Eland's senna
<i>Datura ferox</i>	Large thorn apple
<i>Datura stramonium</i>	Thorn apple
<i>Euphorbia geniculata</i>	Painted milkweed
<i>Sida cordifolia</i>	Heartleaf sida

DIRECTION FOR USE USE ONLY AS DIRECTED

5.1 COMPATIBILITY

The compatibility of **COTTONEX P** products may be influenced by the formulation of the products involved as well as the quality of the water. Since the formulation of other products may change without the knowledge of Makhteshim-Agan and the quality of water may vary from farm to farm, a physical compatibility test should always be carried out prior to application.

5.2 MIXING INSTRUCTIONS

Shake well before use. Replace cap after use.

Half-fill the spray tank with water and pour the required quantity of **COTTONEX P** through a 50 mesh sieve into the tank whilst stirring. Top up the tank with water to the final volume required.

Ensure thorough agitation of the mixture in the tank during mixing and spraying operations.

Tank mixtures must be sprayed out immediately and not allowed to stand in the spray tank.

5.3 APPLICATION TECHNIQUES

COTTONEX P may be applied with any medium or high volume sprayer equipped with an efficient agitation mechanism, which is capable of adequate coverage and even distribution.

For pre-emergence application, best results are obtained using fan-type spray nozzles and applying a spray volume of 200 l water / ha.

When **COTTONEX P** is applied as a directed post-emergence treatment, a floodjet spray nozzle should be used. The use of shields is recommended to prevent the spray mixture reaching the cotton.

5.3.1 Pre-emergence application

COTTONEX P should be applied at or immediately after planting into a freshly cultivated, fine, even, firm and weed free seedbed. To obtain effective weed control it is essential that the application be followed by a light overhead irrigation (10 – 15 mm) to leach the herbicide into the soil before the weeds emerge (within 3 – 4 days after application).

5.3.2 Post-emergence directed application

COTTONEX P may be applied post-emergence as soon as the cotton is tall enough to permit a properly directed spray. Special care must be taken to keep the spray away from the cotton plants as injury may occur. Injury to the bottom leaves does not however, affect yield. Application may take place immediately after an inter-row cultivation or on weeds not more advanced than the 2 – leaf stage. If application takes place immediately after an inter-row cultivation, overhead irrigation should follow to activate the herbicide.

5.4 APPLICATION RATES

COTTON

5.4.1 Pre-emergence application

COTTONEX P is only recommended when used in combination with a suitable grass killer such as **TRIFLUREX 480 EC** (Reg no L5350).

When **COTTONEX P** is used in combination with **TRIFLUREX**, the **TRIFLUREX** must be applied and incorporated according to the manufacturer's recommendations prior to planting the cotton. **COTTONEX P** is then applied as a pre-emergence treatment after the cotton has been planted or during the planting process.

TABLE 1 : **Application rates for COTTONEX P applied pre-emergence or immediately after planting on cotton:**

Soil Type	%Clay	Following pre-plant incorporated grass-killer treatment
		COTTONEX P (1/ ha)
Sand/sandy loam	0 - 20	NOT RECOMMENDED
Sandy clay loam	21 – 30	3,0
Sandy clay loam / sandy clay	31 – 50	4,0
Clay	> 50	NOT RECOMMENDED

Important

- For band treatment calculate the amount of herbicide required in accordance with band width and row spacing. Ensure proper fertilisation to promote vigorous seedling growth.

5.2.4 **Post-emergence directed application**

Apply **COTTONEX P** post-emergence directed as follows, using a suitable surfactant at 0,2% by volume if weeds are present at time of application.

TABLE 2: **Application rates for COTTONEX P applied post-emergence on cotton:**

Soil Type	% Clay	COTOGARD 500 SC (1/ ha)
Sand / sandy clay loam	0 – 30	3,0
Sandy clay loam/ sandy clay	31 – 50	4,0
Clay	> 50	4,0*
Turf	-	4,0*

- On clay and turf soils apply only as directed post-emergence treatment with the weeds in the cotyledon to 2 leaf stage. Pre-emergence applications are **NOT** to be made on such soil types.

COTTONEX P= registered trade mark of Makhteshim-Agan, Ashdod, ISRAEL.

WAARSKUWINGS

- Giftig indien ingesluk.
- Giftig vir visse.
- Berg in 'n koel plek.
- Berg weg van voedsel en voer.
- Hou buite bereik van kinders, oningeligte persone en diere.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid van toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie. Die werking en doeltreffendheid daarvan kan beïnvloed word deur faktore soos abnormale klimaats- en bergingstoestande, kwaliteit van die verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui word nie en die voorkoms van weerstand van die onkruid teen die betrokke middel sowel as die tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid van skade aan gewasse, plantegroei en die omgewing of nadelige invloede op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van versuim van die gebruiker om etiketaanwysings na te kom, of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS

- Moet nie die spuitnewel inasem nie.
- Vermy velkontak.
- Was met seep en water na gebruik.
- Was besoedelde klere na gebruik.
- Moet nie eet, drink of rook tydens vermenging of toediening van die produk of voordat die hande en gesig gewas is nie.
- Moet nie meng en laai nader as 15 m vanaf boorgate, strome, riviere en damme nie.
- Moet nie nader as 15 m van boorgate, strome en riviere toedien nie.
- Moet nie nader as 60 m toedien van damme nie.
- Verseker dat geen terugsyfering plaasvind na boorgate of damme wanneer die produk toegedien word deur 'n besproeiingsstelsel nie.
- Vermy die oorwaai van spuitnewel op ander gewasse, weiveld, riviere, damme, boorgate en plekke wat nie behandel moet word nie.
- Maak toedieningsapparaat skoon na gebruik. Uitspoelwater moet nie gewasse, weiveld, riviere, damme en boorgate besoedel nie.
- Voorkom die besoedeling van voedsel, voer, drinkwater en eetgerei.
- Spoel die leë houer drie keer uit met 'n volume skoon water gelykstaande aan 'n minimum van 10% van die houer. Gooi die spoelwater by die inhoud van die spuittenk voordat die houer vernietig word.
- Moet die leë houer vir geen ander doel gebruik nie.

GEBRUIKSBEPERKINGS

- Moet nie **COTTONEX P** vooropkom toedien op grond waar 'n aanval van saailingverwelksiekte verwag word nie, tensy doeltreffende maatreëls getref is om die siekte bestry te.
- Moet nie **COTTONEX P** vooropkom toedien op gronde wat minder as 20% klei bevat nie.
- Moet nie **COTTONEX P** vooropkom toedien op gruiserige gronde nie.

- Moet geen gewas anders as katoen plant op lande wat met **COTTONEX P** in dieselfde seisoen behandel is nie.
- Moet nie **COTTONEX P** vooropkom op nuutgevormde alluviale gronde soos in die Upington- en Burgersfortgebiede of op die walle van die Limpopo rivier voorkom, gebruik nie.
- Moet nie **COTTONEX P** as 'n na-opkombespuiting gebruik indien dit alreeds vooropkom toegedien is nie.
- Moet nie **COTTONEX P** op brak- en alkalise grond gebruik nie.

ONKRUIDE WAT BEHEER WORD

Die volgende breëblaaronkruid word normaalweg deur **COTTONEX P** beheer wanneer dit in kombinasie met 'n geregistreerde grasdoder gebruik word:

<i>Acanthospermum hispidum</i>	Regopsterklits
<i>Amaranthus hybridus</i>	Gewone misbredie
<i>Amaranthus spinosus</i>	Doringmisbredie
<i>Amaranthus thunbergii</i>	Rooimisbredie
<i>Bidens bipinnata</i>	Spaanse knapsekêrel
<i>Bidens pilosa</i>	Knapsekêrel
<i>Flaveria bidentis</i>	Smelterbossie
<i>Galinsoga parviflora</i>	Knopkruid
<i>Nicandra physaloides</i>	Basterappelliefie
<i>Physalis angulata</i>	Wilde-appelliefie
<i>Portulaca oleracea</i>	Porslein
<i>Richardia brasillensis</i>	Tropiese richardia
<i>Schkuhria pinnata</i>	Kleinkakiebos
<i>Sesbania punicea</i>	Rooi-Sesbania
<i>Solanum nigrum (kompleks)</i>	Nastergal
<i>Tagetes minuta</i>	Kakiebos
<i>Zeleya pentandra</i>	Zeleya
Onkruid wat wisselvallig beheer word	
<i>Cassia italica</i>	Elandsertjie
<i>Datura ferox</i>	Grootstinkblaar
<i>Datura stramonium</i>	Stinkblaar
<i>Euphorbia geniculata</i>	Gekleurde melkkruid
<i>Sida cordifolia</i>	Hartblaartaaiman

GEBRUIKSAANWYSINGS GEBRUIK SLEGS SOOS AANGEDUI

5.1 VERENIGBAARHEID

Die verenigbaarheid van **COTTONEX P** met ander produkte kan beïnvloed word deur die formulاسie van die betrokke produkte asook die kwaliteit van die water. Aangesien die formulاسie van produkte kan verander sonder die medewete van Makhteshim-Agan en die kwaliteit van die water van plaas tot plaas kan varieer, moet 'n fisiese verenigbaarheidstoets altyd uitgevoer word voordat gespuit word.

5.2 MENGINSTRUKSIES

Skud goed voor gebruik. Plaas skroefdeksel terug na gebruik.

Maak die spuittenk halfvol met water en voeg die benodigde hoeveelheid **COTTONEX P** deur 'n 50 maas sif by terwyl geroer word. Vul die tenk verder met water totdat die finale volume verkry word.

Die spuitmengsel moet tydens vermenging en toediening voortdurend in roering gehou word.

Tenkmengsels moet onmiddelik uitgespuit word en nie toegelaat word om in die spuittenk te staan nie.

5.3 TOEDIENINGSTEGNIEKE

COTTONEX P kan met enige medium- of hoëvolume spuittoerusting met 'n doeltreffende roermeganisme toegedien word, mits 'n eweredige bedekking en verspreiding verkry word.

Vir vooropkomtoediening word die beste resultate verkry deur van waaiertipe-spuitpunte gebruik te maak en 'n minimum spuitvolume van 200 l water / ha toe te dien.

Wanneer **COTTONEX P** as 'n gerigte na-opkombespuiting toegedien word, moet 'n "floodjet" spuitpunt gebruik word. Die gebruik van skerms word aanbeveel ten einde te verhoed dat van die spuitmengsel op die katoen beland.

5.3.1 Vooropkomtoediening

COTTONEX P moet tydens of onmiddelik na plant toegedien word aan 'n varsbewerkte, fyn, egalige, ferm en onkruidvrye saadbed. Vir doeltreffende onkruidbeheer is dit noodsaaklik dat toediening gevolg word deur 'n ligte oorhoofse besproeiing (10 – 15 mm) om die onkruidodder in die grond in te loog voordat die onkruid opkom (binne 3 – 4 dae na toediening.)

5.3.2 Na-opkom gerigte toediening

COTTONEX P kan as 'n gerigte na-opkombespuiting toegedien word sodra die katoen groot genoeg is omm dit moontlik te maak. Noukeurige aandag word vereis om te verhoed dat die spuitmengsel nie op die katoenplante beland nie aangesien dit skade kan veroorsaak. Dit het egter geen effek op opbrengs as die onderste blare raakgespuit word nie. Hierdie toediening kan plaasvind onmiddelik na 'n tussenrybewerking of op onkruide wat nie verder as die 2-blaarstadium gevorder is nie. Indien toediening onmiddelik na 'n tussenrybewerking plaasvind, moet sprinkelbesproei word ten einde die onkruidodder te aktiveer.

5.4 TOEDIENINGSHOEVEELHEDE

KATOEN

5.4.1 Vooropkomtoediening

COTTONEX P word slegs in kombinasie met 'n geskikte grasdoder soos **TRIFLUREX 480 EC** (Reg no L5350) aanbeveel.

Wanneer **COTTONEX P** in kombinasie met **TRIFLUREX** gebruik word, moet **TRIFLUREX** voor plant in die grond ingewerk word volgens die vervaardiger se aanbevelings. **COTTONEX P** word dan as 'n vooropkombehandeling tydens of onmiddelik na plant toegedien.

TABEL 1: **Toedieningshoevelhede van COTTONEX P as vooropkombehandeling tydens of onmiddelik na plantproses.**

Grondtipe	% Klei	Volgende op voorplant grasdoder behandeling
Sand/ sandleem	0 – 20	NIE AANBEVEEL
Sandklei leem	21 – 30	3,0
Sandklei leem/ sandklei	31 – 50	4,0
Klei	> 50	NIE AANBEVEEL

Belangrik

- Vir strookbehandeling moet die benodigde hoeveelheid onkruiddoder bereken word volgens die strookbreedte en rywydte.
- Verseker voldoende bemesting om aktiewe saailinggroei te verkry.

5.4.2 **Gerigte na-opkomtoediening**

Dien 'n na-opkombespuiting van **COTTONEX P** toe soos volg en gebruik 'n geskikte benattingsmiddel teen 0,2% per volume indien onkruid teenwoordig is ten tye van toediening.

TABEL 2:

Grondtipe	% Klei	COTTONEX P (l / ha)
Sand / sandkleileem	0 – 30	3,0
Sandkleileem / sandklei	31 – 50	4,0
Klei	> 50	4,0*
Turf	-	4,0*

- * Op klei- en turfgronde dien slegs gerigte na-opkombespuittings toe wanneer die onkruid in die saailing tot 2-blaarstadium is. Vooropkomtoedienings moet **NIE** op sodanige gronde gemaak word nie.

COTTONEX P= geregistreerde handelsmerk van Makhteshim-Agan LTD, Ashdod, ISRAEL.