



FOLPAN 80 WP

Reg No L 6286 Act / Wet 36 of 1947

A wettable powder organic based fungicide for the prevention of anthracnose, downy mildew, deadarm disease and Botrytis in wine grapes, deadarm disease, anthracnose and downy mildew pre-blossom in table grapes as well as late blight and early blight in potatoes.

'n Benatbare poeier organies-gebaseerde swamdoder vir die voorkoming van antraknose, donsige skimmel, streepvleksiekte en Botritis-vrot op wyndruiwe; streepvleksiekte, antraknose en donsige skimmel voorblom op tafeldruiwe asook laat- en vroeë roes op aartappels.

FRAC FUNGICIDE GROUP CODE

M4

FRAC SWAMDODER GROEPKODE

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Folpet (phthalimide)

800 g / kg

Folpet (ptalimied)

NET VOLUME / NETTO VOLUME

..... kg

REGISTRATION HOLDER / REGISTRASIEHOUER

Makhteshim-Agan SA (Pty) LTD

Reg No. 92/01741/07

21 Viben Street

Brackenfell

7560

Batch number

.....

Lotnommer

Date of Manufacture

.....

Datum van Vervaardiging

**FOLPAN is the registered trademark of Makhteshim-Agan Israel.
FOLPAN is die geregistreerde handelsmerk van Makhteshim-Agan, Israel.**

UN NR:



WARNINGS

- Do not use later than pea size on table grapes and on wine grapes not later than 14 days before harvesting.
- Toxic to fish. Store in a cool place away from food and feed. Keep out of reach of children and animals.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will not be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Avoid inhalation of spray mist and dust.
- In case of skin contact wash thoroughly with soap and water.
- Wash contaminated clothing.
- Do not eat, drink or smoke whilst applying or preparing the spray mix, or before washing face and hands.
- Prevent spray drift onto other crops, grazing, rivers, dams and any area not under treatment.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Destroy empty container by perforation and flattening and never use for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

COMPATIBILITY: Folpan 80 WP can be combined with most of the commonly used insecticides and fungicides with the exception of Lime, Lime Sulphur and Bordeaux Mixture.

MIXING AND APPLICATION INSTRUCTIONS: Mix the required quantity in a small amount of water to make a smooth cream, before adding to the recommended amount of water in spray tank. Agitate water in spray tank whilst adding Folpan 80 WP and the spray mixture in spray tank during spray operation.

Spray thoroughly to obtain complete coverage of the plant, including the under-surface of the leaves. It is recommended as a preventive spray.

| CROP AND DISEASE | DOSAGE | REMARKS |
|---|---|---|
| <p>GRAPES Anthrachnose, Deadarm Disease, Botrytis rot</p> <p>Downy mildew</p> | <p>125 g / 100 l water</p> <p>125 g / 100 l water</p> | <p>Spray every two weeks or weekly when weather conditions promote disease development</p> <p>Pre-blossom: First application when shoots are 10 cm long. Repeat at 7 to 10 day intervals depending on weather conditions promoting disease.</p> <p>Flowering to 80 % petal drop: Apply only when flowering period lasts more than two weeks.</p> <p>After flowering: Apply twice or more with 1 to 2 week intervals, depending on weather.</p> |
| <p>POTATOES (In Sandveld area only) Late blight (<i>Phytophthora infestans</i>)</p> <p>Early blight (<i>Alternaria solani</i>)</p> | <p>1.25 kg /ha</p> | <p>Preventative treatment: Apply from emergence at 7 – 10 day intervals in not less than 500 l spray mixture / ha in the pre-blossom stage and in 1 000 l / ha in the post-blossom stage. For the control of early and late blight, particularly after flowering, it is important to obtain thorough coverage.</p> |

WAARSKUWINGS

- Moet nie na ertjiegrootte op tafeldruiwe gebruik nie en op wyndruiwe nie later as 14 dae voor pluktyd nie.
- Giftig vir vis. Stoor in 'n koel plek en weg van voedsel en voer. Hou buite bereik van kinders en diere.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande; kwaliteit van verdunningswater; verenighbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens en dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim deur die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in geval van enige onsekerheid.

VOORSORGMAATREËLS

- Voorkom inaseming van spuitmis en poeier.
- In geval van toevallige velkontak was met seep en water.
- Was besoedelde klere. Moenie eet, drink of rook tydens toediening of vermenging nie, of voordat gesig en hande gewas is nie.
- Voorkom oorwaai van spuitnewel na ander gewasse, weiding, riviere, damme en enige ander gebied wat nie behandel word nie.
- Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.
- Vernietig leë houers deur gate daarin te maak en plat te druk.
- Moet vir geen ander doel gebruik word nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERENIGBAARHIED: Folpan 80 WP kan gemeng word met die meeste insek- en swamdoders wat algemeen gebruik word, met uitsondering van kalk, kalkswael en Bordeaux Mengsel.

MENG EN TOEDIENINGSINSTRUKSIES: Meng die aanbevole hoeveelheid met 'n bietjie water om 'n gladde room te vorm en voeg dan by die res van die hoeveelheid water in spuittenk. Roer water in spuittenk terwyl Folpan 80 WP bygevoeg word en die spuitmengsel in spuittenk tydens bespuiting.

Bespuit deeglik om 'n volledige bedekking te verkry, en sorg veral ook dat die onderkant van die blare deeglik benat word. Dit word aanbeveel as 'n voorkomende bespuiting.

| GEWAS EN SIEKTE | DOSIS | OPMERKING |
|---|--|--|
| DRUIWE Antraknose, Streepvleksierte, Botritisvrot Donsige skimmel | 125 g per 100 l water 125 g per 100 l water | Dien al om die tweede week toe of elke week wanneer weersomstandighede gunstig is vir siekteontwikkeling. Voor bloei: Eerste toediening sodra lote 10 cm lank is. Herhaal met 7 tot 10 dae tussenposes, afhangende van weersomstandighede wat siekte mag begunstig. Bloei tot 80% blomval: Dien slegs toe as bloeitydperk langer as 14 dae duur. Na bloei: Dien twee of meer keer toe met 1 tot 2 weke tussenposes, afhangende van weersomstandighede. |
| AARTAPPELS (Slegs in Sandveldgebied) Laatroes (<i>Phytophthora infestans</i>) Vroeë roes (<i>Alternaria solani</i>) | 1.25 kg / ha | Voorkomende behandeling: Dien toe vanaf opkoms met tussenposes van 7 – 10 dae teen nie minder as 500 l spuitmengsel / ha voor-blom periode en 1 000 l ha in die na-blom stadium. Vir die beheer van vroeë en laatroes, veral na-blom, is dit belangrik om volledige bedekking te verkry. |