

MAKHTESHIM-AGAN SA (PTY) LTD

FOLPAN SULPHUR DP

Reg No L 5454 Act / Wet 36 of 1947

A dusting powder with preventive, fungicidal and miticidal properties for the control of certain fungal diseases and mites as listed on vines and ornamentals.

'n Stuifpoeier met voorkomende swam- en mytdodende eienskappe vir die beheer van sekere swamsiektes en myte soos gelys op wingerd, blomgewasse en sierplante.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Sulphur	900 g / kg	Swael
Folpet (<i>N-trihalomethylthio</i>)	50 g / kg	Folpet (<i>N-trihalometieltio</i>)

NET CONTENT / NETTO INHOUD

..... kg

REGISTRATION HOLDER / REGISTRASIEHOUER

Makhteshim-Agan SA (Pty) LTD

Reg No. 92/01741/07

21 Viben Street

Brackenfell

7560

Batch number	Lotnommer
Date of Manufacture	Datum van Vervaardiging

**FOLPAN SULPHUR DP is the registered trademark of Makhteshim-Agan Israel.
FOLPAN SULPHUR DP is die geregistreeerde handelsmerk van Makhteshim-Agan,
Israel.**

UN NR:



WARNINGS:

DO NOT APPLY WITHIN 3 DAYS OF HARVEST OF TABLE GRAPES, AND WITHIN 4 TO 6 WEEKS OF HARVEST OF WINE GRAPES. During the post-blossom stages larger quantities of dust is required for the control of powdery mildew which, especially on hot days, may result in sulphur burn.

Keep out of reach of children and animals.

Store away from food and feedstuffs, in a cool place.

Keep away from fire and sparks.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, compatibility with other substances not indicated on the label and the occurrence of resistance of the pest or disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

Avoid breathing of dust. Avoid prolonged contact with skin or eyes. In case of eye contact, flush with plenty of water. Wash with soap and water after use. Prevent drift of powder onto other crops, grazing, rivers, dams or any area not under treatment. Destroy empty container and never re-use for any other purpose.

DIRECTIONS OF USE: USE ONLY AS INDICATED

Best results are obtained when plants are dusted when it is cool and windless and when the plants are wet with dew. The application must be repeated after every heavy downpour of rain. Also ensure that a light, even and complete coverage of the plants is obtained.

CROP / PEST	DOSAGE / ha	REMARKS
TABLE AND WINE GRAPES Powdery mildew, Erinose mite, Botrytisrot, Downy mildew	15 - 20 kg	Start dusting, using lower rate when the shoots are 10 - 30 cm long, and repeat shortly after blossoming. Follow up with a third dusting using higher rate 3 - 4 weeks later. Apply a fourth dusting if necessary.
ORNAMENTALS Powdery mildew, Anthracnose and Downy mildew	15 - 30 kg	Dust every 7 - 14 days.

WAARSKUWINGS:

MOET NIE TOEDIEN BINNE 3 DAE VANAF OES IN GEVAL VAN TAFEL DRUIWE, EN 4 TOT 6 WEKE IN DIE GEVAL VAN WYNDRUIWE. Gedurende na-bloom is groter hoeveelhede stuifpoeier nodig om poeieragtige skimmel te beheer en mag dit veral op warm dae tot swaelbrand aanleiding gee.

Hou buite bereik van kinders en diere.

Stoor apart van voedsel en voer, in 'n koel plek.

Hou weg van vuur en vonke.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die plaag of siekte teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die verbruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestand wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS:

Vermynaseming van poeier. Vermynlandurige kontak met vel of oë. In geval van kontak met oë, spoel met baie water uit. Was met seep en water na gebruik. Vermynoorwaai van poeier na ander gewasse, weiding, riviere, damme of gebiede nie onder behandeling nie. Vernietig leë houers en moet nooit hergebruik vir enige ander doel nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

Die beste resultate word verkry wanneer plante bestuif word gedurende koel, windstil tye en wanneer plante nat gedou is. Die bestuiwing moet in alle gevalle na elke sterk bui reën herhaal word. Maak seker dat 'n ligte, egalige en volledige bedekking van die plante verkry word.

GEWAS / PLAAG	DOSIS / ha	OPMERKINGS
TAFEL EN WYNDRUIWE Poeieragtige skimmel, Erinosemyt, Botritisrot, Donsige skimmel	15 - 20 kg	Begin om te bestuif teen laer dosis op 10-30 cm lootlente en herhaal net na blom. Dien 'n derde bestuiwing teen die hoër dosis 3 - 4 weke later toe. Indien nodig dien 'n vierde bestuiwing toe.
SIERPLANTE Poeieragtige skimmel, Antraknose en Donsige skimmel	15 - 30 kg	Bestuif elke 7 - 14 dae.