

MAKHTESHIM-AGAN SA (PTY) LTD

MERPAN® 80 WDG

Reg No L 5456 Act / Wet 36 of 1947

A water dispersible granular fungicide for the control of diseases of crops as listed.

'n Waterdispergeerbare korrel swamdodende vir die beheer van siektes van gewasse soos aangedui.

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Captab

800 g / kg

Kaptab

NET CONTENTS / NETTO INHOUD

..... kg

REGISTRATION HOLDER / REGISTRASIEHOUER

Makhteshim-Agan SA (Pty) LTD

Reg No. 92/01741/07

21 Viben Street

Brackenfell

7560

Batch number

.....

Lotnommer

Date of Manufacture

.....

Datum van Vervaardiging

**MERPAN 80 WDG is the registered trademark of Makhteshim-Agan Israel.
MERPAN 80 WDG is die geregistreerde handelsmerk van Makhteshim-Agan, Israel.**

UN NR:
(green / groen)



WARNINGS:

Do not harvest within 14 days of last application except in the case of table grapes which is 4-6 weeks and tomatoes 7 days. Keep out of reach of children, uninformed persons and animals. Store in a cool place away from food and feed.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will not be affected by factors such as abnormal climatic and storage conditions; quality of dilution water; compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

Wash hands and all exposed parts of the body thoroughly after use. Avoid inhalation of powder and spray mist. Prevent contamination of food, eating utensils and drinking water. Destroy empty container and do no re-use for any other purpose.

DIRECTIONS OF USE: USE ONLY AS INDICATED**COMPATIBILITY:**

Compatible with non-alkaline materials such as Parathion.

APPLICATION INSTRUCTIONS:

Unless otherwise stated spray at 7 - 10 day intervals and after rain.

CROP / PEST	DOSAGE / HA	REMARKS
POTATOES Late blight	1,25 - 2 kg / 300 l water / ha as a low volume spray 1,25 - 2 kg / 1 000 l water / ha as a high volume spray	Start spraying before disease is due to appear. Use the higher dosage rate for big plants and / or when infection is severe.
TOMATOES Late blight and <i>Septoria</i> leaf spot	125 g / 100 l water	Seedbeds: Spray twice a week. Lands: Start to spray before disease is due to appear. Depending on plant size apply 500 - 2 000 l spray mixture / ha.
APPLES AND PEARS Scab	125 g / 100 l water	Pre-blossom: Commence at green tip stage and repeat at 7 day intervals. Post-blossom: Continue with 3-4 sprays according to weather conditions at 10-14 day intervals.
NOTE: MERPAN WGD may cause leafspotting of Pearmain, Delicious, Cox and other sensitive apple cultivars.		
PEACHES Freckle	125 g / 100 l water	Apply at full blossom, 2 weeks later and 3 weeks after second spray.
TABLE GRAPES Anthracnose	125 g / 100 l water	Apply to young regrowth and bunches when berries start swelling and become translucent. Repeat at

			10 - 14 day intervals. Do not use on wine grapes.
ROSES Black spot	125 g / 100 l water		Start spraying at first signs of disease and repeat every 7 - 10 days.
TURF Fairy ring, Dollar spot and other leaf, crown and root diseases	200 g / 100 l water / 50 m ⁵		Spray at first sign of disease.

WAARSKUWINGS:

Moenie oes binne 14 dae nadat laaste toediening plaasgevind het nie, behalwe in die geval van tafeldruiwe wat 4-6 weke is en tamaties 7 dae. Hou buite bereik van kinders, oningeligte persone en diere. Stoor in 'n koel plek weg van voedsel en veevoer.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die plaag of siekte teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die verbruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestand wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS:

Was hande en alle blootgestelde gedeeltes van die liggaam deeglik na gebruik. Voorkom inaseming van stof of spuitnewel. Voorkom kontaminasie van voedsel, eetgereedskap en drinkwater. Vernietig leë houers en moenie vir enige ander doeleindes hergebruik nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERENIGBAARHEID:

Verenigbaar met nie-alkaliëse produkte soos PARATION.

TOEDIENINGS VOORSKRIFTE:

Tensy anders aanbeveel, wend aan op 7 - 10 dae tussenposes en na reën.



GEWAS / PLAAG	DOSIS / HA	OPMERKINGS
AARTAPPELS Laatroes	1,25 - 2 kg / 300 l water / ha as 'n lae volume toediening of 1,25 - 2 kg / 1000 l water / ha as 'n hoë volume toediening	Begin met toedienings voordat die siekte verskyn Gebruik die hoë dosis vir groot plante of tydens hoë infeksie.
TAMATIES Laatroes en blaarvlek	125 g / 100 l water	Saadbeddings: Spuit tweekeer 'n week. Lande: Begin om te spuit voordat siekte verskyn. Afhangende van plantgrootte dien 500 - 2 000 lt spuitmengsel / ha toe.
APPELS EN PERE Skurf	125 g / 100 l water	Voor Blom. Begin by groenpuntstadium en herhaal met 7 dae tussenposes. Na-blom. Gaan voort met 'n verdere 3 - 4 toedienings afhangende van weerstoestand met 10 - 14 dae tussenposes.
NOTA: MERPAN WDG mag blaarvleke op appel kultivars soos Pearmain, Delicious, Cox en ander sensitiewe appel kultivars veroorsaak.		
PERSKES Sproetsiekte	125 g / 100 l water	Dien toe op volblom, 2 weke later en 3 weke na tweede spuit.
TAFELDRUIWE Antraknose	125 g / 100 l water	Dien toe aan jong lote en trosse wanneer druiwekorrels begin swel en deurskynend word. Herhaal met tussenposes van 10 - 14 dae. Moenie op wyndruiwe gebruik nie.

<p>ROSE Swartvlek</p>	<p>125 g / 100 l water</p>	<p>Begin wanneer eerste tekens van siekte verskyn. Herhaal elke 7 - 10 dae.</p>
<p>GRASPERKE Heksering, Dollar-vlek en ander blaar-, kroon-en- wortelsiektes</p>	<p>200 g / 100 l water / 50m5</p>	<p>Spuit sodra die eerste tekens van die siekte opgemerk word.</p>