



GROUP H FUNGICIDE

GROEP H SWAMDODER

## NIMROD

Reg. No. L 1117 : Act / Wet 36/1947

An emulsifiable concentrate systemic fungicide for the control of powdery mildew in apples, peaches, cucurbits and mangoes.

'n Emulgeerbare konsentraat sistemiese swamdoder vir die beheer van poeieragtige skimmel in appels, perskes, pampoengewasse en mango's.

**ACTIVE INGREDIENT:**

bupirimate (pyrimidine)

250 g / l

**AKTIEWE BESTANDDEEL:**

bupirimaat (pirimidien)

NET VOLUME / NETTO VOLUME

.....

Registration holder / Registrasiehouer:

**MAKHTESHIM-AGAN SA (PTY) LTD.**

Reg. No. 92/01741/07

21 Viben Street

BRACKENFELL 7560

Tel (021) 982 1460

Fax (021) 982 5810

Batch number..... Lotnommer

Date of manufacture..... Datum van vervaardiging

**BEFORE YOU OPEN THE CONTAINER READ THE ENTIRE LABEL  
LEES DIE ETIKET VOLLEDIG VOORDAT U DIE HOUER OOPMAAK**

UN No. 1993

VN No. 1993



**HARMFUL  
SKADELIK**



CLASS II  
band

Yellow

Nimrod

**WARNINGS:**

- \* **Withholding periods:** Minimum number of days between last application and harvest.

**Apples, Peaches and Nectarines** ..... 14  
days

**Cucurbits:** .....  
.....3 days

- \* Harmful when swallowed.
- \* Irritating to eyes and skin.
- \* May cause sensitisation by skin contact.
- \* Toxic to fish.
- \* **Flammable** - keep away from open flames.
- \* Store in a cool place away from food and feed.
- \* Keep out of reach of children, uninformed persons and animals.
- \* In case of poisoning call a doctor and make this label available to him.

- \* **General warning:**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions; quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

**PRECAUTIONS:**

- Do not eat drink or smoke whilst handling this product.
- Prevent contamination of food, feed, drinking water and eating utensils.
- Wear rubber gloves and face shield when handling the concentrate. If you get it in your eyes, flush it out at once. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them.

- Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsings to the contents of the spray tank before destroying the container.
- While spraying, avoid contact with the spray as much as you can. Avoid spray drift onto other crops, grazing rivers or dams.
- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams. Destroy empty container and do not use for any other purpose. Change and wash your work clothes. Wash yourself.

#### Symptoms of human poisoning:

Incontinence and subdued behaviour with salivation and an increased respiration rate. Massive doses may result in renal failure.

#### First aid and medical treatment:

**Eye contamination:** Hold eyelids apart and pour in a gentle stream of water for 10 - 15 minutes. Go to a doctor.

**If NIMROD is swallowed, do not make the person vomit.** Take person and this product label to a doctor at once for medical treatment.

**Medical treatment:** If gastric lavage is performed, take care to prevent aspiration of gastric contents. Consider administration of activated charcoal and a laxative. Treat symptomatically.

**DIRECTIONS FOR USE:** Use only as indicated.

CROP	DOSAGE	REMARKS
<b>Apples, Peaches and Nectarines</b> Powdery mildew	60 ml / 100 l water	Under conditions of a high infection pressure use the higher concentration. Repeat application at 10 - 14 day intervals.
<b>Cucurbits</b> Powdery mildew	40 - 60 ml / 100 l water	Under conditions of a high infection pressure use the higher concentration. Repeat application at 10 - 14 day intervals.
<b>Mangoes</b> Powdery mildew	40 ml / 100 l water	Apply at 14 day intervals from first signs of infection up to 100 % petal fall. The first application is usually made when 50 % of the flowers on the panicle are open. A further 2 - 3 applications are required depending on

		the weather conditions and uniformity of the flowering period. A high volume application must be made to ensure a good coverage of the flower panicle.
--	--	--

### WAARSKUWINGS:

- \* **Onthoudingsperiodes:** Minimum aantal dae tussen laaste toediening en oes:

**Appels, Perskes en Nektariens:** ..... 14 dae

**Pampoengewasse:** ..... 3 dae

- \* Skadelik indien ingesluk.
- \* Irriterend vir die oë en vel.
- \* Mag sensitisering veroorsaak deur velkontak.
- \* Giftig vir visse.
- \* **Vlambaar** - hou weg van oop vlamme.
- \* Berg in 'n koel plek weg van voedsel en voer.
- \* Hou buite bereik van kinders, oningeligte persone en diere.
- \* In geval van vergiftiging ontbied 'n geneesheer en toon hierdie etiket aan hom.

- \* **Algemene waarskuwing:**

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale grond-, klimaats- en bergingstoestande; kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die siekte teen die betrokke middel, sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

**VOORSORGMATREËLS:**

- Moet nie eet, drink of rook terwyl die produk hanteer word nie.
- Vermoed besoedeling van voedsel, voer, drinkwater en eetgerei.
- Dra rubberhandskoene en gesigs-kerm wanneer die konsentraat hanteer word. Indien dit in u oë kom, spoel onmiddellik uit. Indien dit op u vel kom, was dit dadelik af. Indien dit op u klere mors, trek uit en was dit.
- Keer die leë houër om oor die spuit- of mengtenk en dreineer vir minstens 30 sekondes nadat die vloei tot 'n gedrup verminder het. Spoel die houër daarna driekeer uit met 'n volume water gelykstaande aan 'n minimum van 10 % van die inhoud van die houër. Gooi die spoelwater by die inhoud van die spuittenk voordat die houër vernietig word.
- Vermoed kontak met die spuitstof sover moontlik tydens toediening. Vermoed oorwaai van spuitnewel na ander gewasse, weiding, riviere of damme.
- Maak toediener deeglik skoon na gebruik en gooi waswater waar dit nie gewasse, weiding, riviere of damme sal besoedel nie. Vernietig leë houër en moet dit nie vir enige ander doel gebruik nie. Trek werksklere uit en was dit. Was uself.

**Simptome van menslike vergiftiging:**

Onbeheertheid en stil gedrag met kwyling en 'n toename in asemhalingspoed. Groot dosisse mag akute nierversaking tot gevolg hê.

**Noodhulp- en mediese behandeling:**

**Oogbesoedeling:** Hou ooglede oop en spoel met 'n ligte stroom water vir 10 - 15 minute. Gaan na 'n dokter.

**Indien NIMROD ingesluk is, moet nie braking veroorsaak nie.**

Neem die persoon en hierdie produk-etiket onmiddellik na 'n dokter vir mediese behandeling.

**Mediese behandeling:** Indien 'n maagspoeling gedoen word, voorkom aspirasie van die maaginhoud. Oorweeg toediening van geaktiveerde houtskool en 'n purgeermiddel. Behandel simptome.

**GEBRUIKSAANWYSINGS:** Gebruik slegs soos aangedui.

GEWAS	DOSIS	OPMERKINGS
Appels, Perskes en Nektariens Poeieragtige	60 ml / 100 l water	Onder toestande van 'n hoë besmettingsdruk gebruik die hoër konsentrasie. Herhaal toediening elke 10 - 14 dae.

skimmel		
<b>Pampoengewasse</b> Poeieragtige skimmel	40 - 60 ml / 100 l water	Onder toestande van 'n hoë besmettingsdruk gebruik die hoër konsentrasie. Herhaal toediening elke 10 - 14 dae.
<b>Mango's</b> Poeieragtige skimmel	40 ml / 100 l water	Dien toe met 14 dae tussenposes vanaf eerste tekens van besmetting tot en met 100 % blomblaarval. Die eerste toediening geskied gewoonlik wanneer 50 % van die blomme op die blomsteel oop is. 'n Verdere 2 - 3 bespuitings moet toegedien word afhangede van die weers-toestande en egaligheid van die blomperiode. Dien 'n hoëvolume bespuiting toe om 'n goeie bedekking van die blomtrosse te verseker.